

# Ectoparasites and Pathogens of Kuhl's Pipistrelle *Pipistrellus kuhlii* (Kuhl, 1817) (Chiroptera: Vespertilionidae): Our Own and Published Data Review

M. V. Orlova<sup>a, b, \*</sup>, D. G. Smirnov<sup>c, \*\*</sup>, V. P. Vekhnik<sup>d, \*\*\*</sup>,  
A. M. Lukyanenko<sup>c, \*\*</sup>, and A. V. Zabashta<sup>e, \*\*\*\*</sup>

<sup>a</sup>Tyumen State University, Tyumen, 625003 Russia

<sup>b</sup>Tomsk National Research State University, Tomsk, 634050 Russia

<sup>c</sup>Penza State University, Penza, 440026 Russia

<sup>d</sup>Spryginsk State Natural Biosphere Reserve, Zhigulevsk, 445362 Russia

<sup>e</sup>Rostov-on-Don Anti-Plague Scientific Research Institute, Rostov-on-Don, 344002 Russia

\*e-mail: masha\_orlova@mail.ru

\*\*e-mail: eptesicus@mail.ru

\*\*\*e-mail: vekhnik@mail.ru

\*\*\*\*e-mail: zabashta68@mail.ru

Received December 26, 2019; revised July 21, 2020; accepted August 20, 2020

**Abstract**—Here we report the results of our own survey and literary published data on the ectoparasite fauna and pathogens of the alien bat species, the Kuhl's pipistrelle *Pipistrellus kuhlii* (Kuhl, 1817) (Chiroptera: Vespertilionidae). This bat is a host of 36 species of parasitic mites, ticks and insects (including accidental findings) and 13 species of pathogens (protozoa, bacteria, viruses). The flea *Ischnopsyllus variabilis* is recorded on this host for the first time. We have found that outside of the host ancestral range, the core of the bat parasite fauna is significantly different due to the loss of host species-specific ectoparasites. Particularly, in Russia, only 6 species of parasitic arthropods have been recorded for Kuhl's pipistrelle and all of them are host genus-specific. At the same time, the features of ecology and occasional finds of extrinsic parasites allow to suggest that *P. kuhlii* has wide contacts with animals which are the reservoirs of zoonotic infections, that in combination with the fact of isolation of several pathogens from this species (including two coronaviruses) points to a possible medical importance of Kuhl's pipistrelle.

**Keywords:** Kuhl's pipistrelle, *Pipistrellus kuhlii*, bat ectoparasites, *Steatonyssus periblepharus*, *Ischnopsyllus variabilis*, *Carios vespertilionis*

**DOI:** 10.1134/S2075111720040104

## INTRODUCTION

The Kuhl's pipistrelle or Mediterranean *Pipistrellus kuhlii* (Kuhl 1817), originating from the south (Strelkov and Il'in, 1990; Vernier and Bogdanowicz, 1999), originally preferred open plain arid and semi-arid landscapes of the Mediterranean, North Africa, Near East, and Kashmir (Strelkov et al., 1985). Progressive spread of this species was reported over the last several decades in Europe (Il'in, 2000; Strelkov, 2004; Bogdanowicz, 2004; Sachanowicz et al., 2006). This contributed to the expansion of the total area of the range by a factor of five (Ancillotto et al., 2016). A pronounced tendency toward synanthropy, high adaptability to various climatic conditions, and general global warming induced its rapid dispersal in the northerly and easterly directions. From the direction of the south of Central and Eastern European, the spe-

cies expansion northward was first reported as early as the 1980s, when *P. kuhlii* was recorded from the north of the Alps in Switzerland, southern Germany, Austria, and Hungary (Haffner et al., 1991; Feher, 1995; Bauer, 1996; Meschede et al., 1998; Fiedler et al., 1999; Spitzerberger and Bauer, 2001). Currently, the species has reached the latitude of 50° and was reported from Slovakia (Ceeľuch and Ševčík, 2006; Danko, 2007), the Czech Republic (Reiter et al., 2007), and Poland (Sachnowicz et al., 2006). Additionally, the active dispersal was recorded in Ukraine (Kedrov and Shurshak, 1999; Godlevsky et al., 2000; Poluda and Zagorodnyuk, 2001; Gavris' and Kotserzhinskaya, 2002; Zagorodnyuk and Reznik, 2007; Gashchak et al., 2009; Godlevs'ka, 2015), Belarus (Demyanchik, 2013; Shpak and Larchenko, 2016), and Central Asia (Strelkov, 2004).

Prior to the mid-20th century, finds of *P. kuhlii* were known to be rare from the regions neighboring Russia, specifically, a few settlements in Transcaucasia and Central Asia and in Crimea (Kuzyakin, 1950; Strelkov, 1963). In the east of European Russia, the starting point of this species expansion was southeastern Transcaucasia, where a rapid buildup of its population level was observed in the 1940s–1950s (Vereshchagin, 1959). The first finds were made in Rostov-on-Don in 1975 (Yarmysh and Kazakov, 1977), in Grozny and Abrau-Dyurso in Krasnodar territory in 1977 (Yarmysh et al., 1980), and North Ossetia in 1978 (Komarov and Kuchiev, 1982). In the Volga region, *P. kuhlii* was first discovered in 1980 in the vicinity of Lake Baskunchak in Astrakhan region ( $48^{\circ}$  N) (Lindeman and Subbotin, 1983), where the typical plain species was able to arrive exclusively from the south, while migrating along the western coast of the Caspian Sea (Strelkov et al., 1985). Taking into account, however, its subsequent spread northward, it is not implausible that the species could have found its way to the region much earlier. By 1985, its finds had already been reported from the north of Volgograd region, where it reached  $50^{\circ}$  N (Strelkov et al., 1985). *P. kuhlii* was detected in Saratov region at  $51^{\circ}$  latitude in 1988 (Strelkov and Il'in, 1990) and reported from  $52^{\circ}$  N as early as 1995 (Zav'yalov and Shlyakhtin, 1999). In 1997, the animals were already found by us in Samara, whereas soon after, its habitat was identified in the cities of Novokuibyshevsk and Tolyatti (Smirnov and Vekhnik, 2011). The species was found at Samara Bend and in Penza and Ul'yanovsk regions in the early 2000s (Il'in et al., 2006; Zolina et al., 2007; Smirnov et al., 2007; Shepelev et al., 2008; Smirnov and Vekhnik, 2011), as well as in the Republic of Mordovia (Artayev and Smirnov, 2016). About the same time, it was detected in Voronezh (Smirnov and Vekhnik, 2011), Tambov (Lada, 2010), Bryansk (Sitnikova et al., 2009), and Moscow (Kruskop and Kozhurina, personal communication) regions. Prior to 1990 in the Transvolga, finds of *P. kuhlii* were limited to occasional spots located easterly up to the Bolshoy Uzen River within the Volga-Ural interfluvium (Strelkov and Il'in, 1990). Further eastward, the species was found at the end of the 20th century strictly in the proximity to the Ural River Delta (Shaimardanov, 2001). Attempts made in the 1990s and in the early 2000s with intent to find it in the floodplain of the lower and middle reaches of the Ural River yielded no result; but in 2008, *P. kuhlii* was detected at a number of points in Orenburg and Chelyabinsk regions nevertheless (Davygora et al., 2009). The northmost find was made in Udmurtiya (Izhevsk) in winter 1991, but the animal was probably transported there accidentally by vehicle from the south (Kapitonov and Grigor'ev, 1995).

Despite the spotlight recently put on the Kuhl's pipistrelle (Smirnov and Vekhnik, 2011; Godlevs'ka, 2015; Shpak and Larchenko, 2016; Ancillotto et al., 2016; Sachnowicz et al., 2017), the parasitic fauna and

pathogens associated with the species are considered in scattered fragmentary communications from various (primarily, African, Asian, and South European) parts of the range. As for Russia, five or six species of ectoparasites were described earlier. No information, in actual fact, is available from Western Europe. From the epidemiological perspective, bats are, however, known to be a huge reservoir of zoonotic infections. The literature mentions at least 250 virus agents alone, including infectious agents of dangerous diseases, such as rabies, Ebola fever, SARS, MERS, and COVID-2019 isolated from microbats or otherwise detected in their body (Schountz, 2013; Moratelli and Calisher, 2015), while, in addition, many of their ectoparasites are involved in circulation of agents of different nature (Orlova and Kononova, 2018). As already been mentioned, the Kuhl's pipistrelle demonstrates active expansion across the territory of Russia and, therefore, may possibly transport the temporal ectoparasites over a considerable distance, thus causing a number of diseases to spread further.

## MATERIALS AND METHODS

Bats were captured using the trapping nets during the period of 2006–2019; the age and sex were determined in each individual and the reproductive state (pregnancy and lactation) was determined in females (Racey, 2009). Thirty-nine individuals of the studied species were captured in Astrakhan region by D.G. Smirnov, V.P. Vekhnik, and A.M. Luk'yanenko in the following localities: village of Ivanchug ( $46^{\circ}04'$  N,  $47^{\circ}59'$  E; May 22, 2018, four specimens); Astrakhan Nature Reserve, Damchikskii site ( $45^{\circ}47'$  N,  $47^{\circ}53'$  E; May 23, 2018, 17 specimens); Astrakhan Nature Reserve, Obzhorovskii cordon (guard station) ( $46^{\circ}18'$  N,  $48^{\circ}59'$  E; August 27 and 28, 2019; 18 specimens). In Dagestan, pipistrelles were captured by Smirnov and Vekhnik in the village of Kochubei ( $44^{\circ}24'$  N,  $46^{\circ}32'$  E; April 24, 2019; five individuals examined), in the outskirts of Khasavyurt on the Aktash River ( $43^{\circ}17'$  N,  $46^{\circ}38'$  E; May 20, 2019; three specimens), and in the Agrakhanskii Sanctuary of the Dagestan State Natural Reserve ( $43^{\circ}48'$  N,  $47^{\circ}31'$  E; May 23, 2019; three specimens). One animal of this species was captured by Smirnov in the village of Novospasskoe, Ul'yanovsk region ( $53^{\circ}08'$  N,  $43^{\circ}45'$  E; August 17, 2006). In Rostov region, one individual was trapped by Zabashta in the outskirts of Rostov-on-Don ( $47^{\circ}29'$  N,  $39^{\circ}56'$  E; July 20, 2019). Thus, a total of 52 individuals of *P. kuhlii* were examined in eight localities of the Russian south; after the examination and collection of ectoparasites, all bats were released in the wild.

Ectoparasites were sampled using tweezers and needle and fixed in 70% ethanol. Permanent preparations were made using Faure-Berlese mounting medium according to standard technique (Whitaker, 1988); parasites were identified by M.V. Orlova using standard keys and other taxonomic publications (Hopkins

**Table 1.** Distribution of the Kuhl's pipistrelle across the localities (*n* is the number of examined hosts; first line is the absolute number; second line is MI; and third line is *P*, %)

Locality	Astrakhan region			Dagestan			Rostov region	Ul'yanovsk region	Total			
	Astrakhan Nature Reserve		Ivanchug village	Kochubei village	Khasavyurt outskirts	Agrakhanskii Sanctuary						
	Obzhorovskii cordon	Damchikskii site										
Species of parasite	<i>n</i> = 18	<i>n</i> = 17	<i>n</i> = 4	<i>n</i> = 5	<i>n</i> = 3	<i>n</i> = 3	<i>n</i> = 1	<i>n</i> = 1	<i>n</i> = 52			
<i>Steatonyssus periblepharus</i>	10 1 56	—	—	18 3.6 100	7 3.5 67	10 3.3 100	7	—	52 2.5 40			
<i>Carios vespertilionis</i>	4 2 11	—	—	—	—	—	—	—	4 2 4			
<i>Ischnopsyllus octactenus</i>	5 1 28	3 1 18	5 1.3 100	2 2 20	—	1 1 33	—	—	16 1.14 27			
<i>Ischnopsyllus variabilis</i>	—	—	—	—	—	—	—	1	1 1 2			
Total	19 2.1 50	3 1 18	5 1.3 100	20 4 100	7 3.5 67	11 3.7 100	7 1	1	73 2.6 54			

and Rothschild, 1956; Filippova, 1966; Orlova et al., 2016). The core of the parasitic fauna was assessed on the basis of the parameters proposed by Balashov (2009). Host infestation parameters are presented using the standard parasitology indices, such as MI (mean intensity), which is the mean number of parasites per infested host, and *P* (prevalence), which is percentage of infested individuals.

## RESULTS

There were 73 specimens of four ectoparasite species (in Acari and insects) collected from the examined individuals of *P. kuhlii* (Table 1).

### Acari: Gamasina: Macronyssidae

*Steatonyssus periblepharus* Kolenati, 1858: 4 ♀♀, 6 N1 (Astrakhan Nature Reserve, Obzhorovskii cordon, August 27 and 28, 2019); 13 ♀♀, 5 N1 (Dagestan, Kochubei village, April 24, 2019); 3 ♀♀, 4 N1 (Dagestan, Khasavyurt outskirts, May 20, 2019); 9 ♀♀ (including one with intrauterine egg); N1 (Dagestan, Agrakhanskii Sanctuary, May 23, 2019); ♂, 6 N1 (Rostov region, Platov Airport, July 20, 2019).

### Acari: Parasitiformes: Argasidae

*Carios vespertilionis* Latreille, 1796: 4 L (Astrakhan Nature Reserve, Obzhorovskii cordon, August 27 and 28, 2019).

### Insecta: Siphonaptera: Ischnopsyllidae

*Ischnopsyllus octactenus* (Kolenati, 1856): 5 ♀♀ (Astrakhan region, Ivanchug, May 22, 2018); 2 ♀♀, ♂ (Astrakhan Nature Reserve, Damchikskii site, May 23, 2018); 4 ♀♀, ♂ (Astrakhan Nature Reserve, Obzhorovskii cordon, August 27 and 28, 2019); ♀ and ♂ (Dagestan, Kochubei village, April 24, 2019); ♀ (Dagestan, Agrakhanskii Sanctuary, May 23, 2019).

*Ischnopsyllus variabilis* (Wagner, 1898): ♀ (Ul'yanovsk region, Novospasskoe village, August 17, 2006).

## DISCUSSION

Our samples feature the dominant number of *S. periblepharus* (52 out of 73 specimens (in other words, nearly three-fourth of the collected parasites) found in five out of eight inspected localities. The intensity of their infestation of bats is relatively low (MI not exceeding 3.6); the proportion of infested

**Table 2.** Anthropods collected from the Kuhl's pipistrelle and their medical importance

Ectoparasite	Finds in Russia	Finds outside of Russia	Principal hosts	Medical importance
Acar: Prostigmata: Myobiidae				
<i>Pieracarus pipistrellus pipistrellus</i> (Radford, 1938)	—	Armenia (Dusbabek and Arutunian, 1976)	Pipistrelles <i>Pipistrellus</i> spp.	—
<i>Acanthophthirius kolenatii</i> Dusbabek, Arutunian, 1976	—	Armenia (Dusbabek and Arutunian, 1976)	Type host is <i>P. kuhlii</i>	—
<i>Calcaromyobia</i> sp.	—	Azerbaijan (Dubovchenko, 1968)	—	—
<i>Myobiidae</i> gen. sp.	—	Azerbaijan (Dubovchenko, 1968)	—	—
Acar: Prostigmata: Trombiculidae				
<i>Oudemansidium komareki</i> (Daniel et Dusbabek, 1959)	—	Spain (Minorca Island) (Stekolnikov and Quetglas, 2019)	Multiple horseshoe and simple nosed microbats; finds from rodents are known	—
Acar: Gamasina: Ameroseiidae				
<i>Kleemannia</i> sp.	—	Azerbaijan (Dubovchenko, 1968)	Mites of this genus do not parasitize	—
Acar: Gamasina: Laelaptidae				
<i>Laelaps algericus</i> Hirst, 1925	—	Azerbaijan (Dubovchenko, 1968)	Rodents (Rodentia), more commonly house mouse <i>Mus musculus</i> Linnaeus, 1758	Carrier of plague <i>Yersinia pestis</i> (Lehmann and Neumann 1896) van Loghem 1944 and lymphocytic choriomeningitis virus LCMV (Arenaviridae) (Zemskaya, 1973)
Acar: Gamasina: Dermanyssidae				
<i>Dermanyssis gallinae</i> Redi, 1674	—	Azerbaijan (Dubovchenko, 1968)	Domestic and wild birds (Aves)	Cause dermatosis; carry paramyxoviridae, horse encephalomyelitis virus, birds' causative agents <i>Pasteurella multocida</i> (Lehmann and Neumann 1899) Rosenbusch and Merchant 1939, <i>Salmonella gallinarum</i> (Flochlay et al., 2017)
Acar: Gamasina: Spiturnicidae				
<i>Meristaspis lateralis</i> Kolenati, 1857	—	Iran (Benda et al., 2012)	Megabats from the family Pteropodidae (Chiroptera: Pteropodidae)	—

Table 2. (Contd.)

Ectoparasite	Finds in Russia	Finds outside of Russia	Principal hosts	Medical importance
<i>Spinturnix acuminatus</i> (Koch, 1836)	—	Palestine (Anciaux de Faveaux, 1976)	Noctule bats <i>Nyctalus</i> spp.	—
<i>Spinturnix acuminata</i> group	—	Libya (Benda et al., 2014)	—	—
<i>Spinturnix bakeri</i> Rudnick, 1960	—	Azerbaijan (Dubovchenko, 1968)		—
<i>Spinturnix myoti</i> (Kolenati, 1856)	—	Azerbaijan (Gadzhiev and Dubovchenko, 1967)	Mouse-eared bat <i>Myotis</i> spp.	Carry bacteria <i>Bartonella</i> sp. (Hornok et al., 2012)
<i>Spinturnix kolenatii</i> Oudemans, 1910	—	Azerbaijan (Gadzhiev and Dubovchenko, 1967)	<i>Eptesicus</i> spp.	—
<i>Spinturnix psi</i> (Kolenati, 1856)	—	Armenia (Arutyunyan and Ogadzhanyan, 1974)	Long-winged bats (Miniopteridae)	Carry bacteria <i>Anaplasma</i> <i>phagocytophillum</i> (Foggie 1949) Dumler et al., 2001 (Reeves et al., 2006)
Acarai: Gamasina: Macronyssidae				
<i>Macronyssus cyclaspis</i> (Oudemans, 1906)	The former Soviet Union, no indication of specific state (Stanyukovich, 1997)		Pipistrelle <i>Pipistrellus</i> spp.	—
<i>Macronyssus flavus</i> (Kolenati, 1856)	—	Azerbaijan (Dubovchenko, 1968)	Noctule bats <i>Nyctalus</i> spp.	—
<i>Macronyssus kolenatii</i> (Oudemans, 1902)	Rostov region (Zabashta et al., 2019)	Egypt (Radovsky, 1967)	Pipistrelles <i>Pipistrellus</i> spp.	—
	The former Soviet Union, no indication of specific state (Stanyukovich, 1997)			
	Rostov region (Zabashta et al., 2019; present article), Astrakhan region (Zabashta et al., 2019; present article), Dagestan (present article)	Azerbaijan (Dubovchenko, 1968); Armenia (Ogadzhanyan and Arutyunyan, 1974); Israel (Korine et al., 2017); Jordan (Benda et al., 2010); Palestine (Anciaux de Faveaux, 1976); Iran (Benda et al., 2012); Lybia (Benda et al., 2014)	Pipistrelles <i>Pipistrellus</i> spp.	Carry bacteria <i>Borellia afzelii</i> Canica et al., 1994 (Zabashta et al., 2019)
<i>Steatonyssus periblepharus</i> Kolenati, 1858	—	Iran (Sharifi et al., 2008)	—	—
<i>Steatonyssus</i> sp.	—	Iran (Sharifi et al., 2008)	—	—

Table 2. (Contd.)

## ECTOPARASITES AND PATHOGENS OF KUHL'S PIPISTRELLE

353

Ectoparasite	Finds in Russia	Finds outside of Russia	Principal hosts	Medical importance
<i>Parasteatonyssus ca. nyctinomi</i>	—	Israel (Korine et al., 2017)	Free-tailed bats <i>Tadarida</i> spp. (Chiroptera: Molossidae)	—
<i>Carios vespertilionis</i> Latreille, 1796	Astrakhan region (Zabashta et al., 2019; present article)	Spain (Imaz et al., 1999) Iran (Sharifi et al., 2008) Israel (Korine et al., 2017) Jordan (Benda et al., 2010) Algeria (Bendjedou et al., 2017) Lybia (Benda et al., 2014)	Pipistrelles <i>Pipistrellus</i> spp.	Carry bacteria <i>Rickettsia</i> sp. AvBat, <i>Ehrlichia</i> sp. AvBat, <i>Borrelia</i> sp. CPB1 (Socolovschi et al., 2012), <i>Babesia vesperuginis</i> Dionisi, 1899 (Apicomplexa: Piroplasmida) (Gardner and Molynieux, 1988; Hornok et al., 2016, 2017); Issyk-Kul Virus ISKV, Keterah orthonaairovirus KTRV (Bunyavirales: Nairoviridae) (Lvov et al., 1973; International Catalogue of Arboviruses..., 1985; Al'khovskii et al., 2013), virus from the fam. Bunyaviridae (similar to Issyk-Kul virus) (Oba et al., 2016), tick-borne encephalitis virus (Flaviviridae) (I.V. Kuz'min and V.V. Yakimenko, personal communication)
<i>Hyalomma dromedarii</i> Koch 1844	—	Algeria (Bendjedou et al., 2017)	Acari: Parasitiformes: Ixodidae Camels (Artiodactyla: Camelidae)	Carrier of protozoa <i>Theileria camelensis</i> (Apicomplexa: Piroplasmida) (Hoogstraal, 1954; Abd El-Baky, 2001; Hamed et al., 2011)
<i>Nycteribia schmidlii</i> Schiner, 1853	—	Turkey (Aktas and Hasbenli 1994)	Long-winged bats <i>Miniopterus</i> spp.	Carrier of virus from the family Rhabdoviridae (Aznar-Lopez et al., 2013)
<i>Nycteribia vexata</i> Westwood, 1835	—	Turkey (Aktas and Hasbenli, 1994)	Mouse-eared bats	—

Table 2. (Contd.)

Ectoparasite	Finds in Russia	Finds outside of Russia	Principal hosts	Medical importance
<i>Penicillidia conspicua</i> Speiser, 1901	—	Turkey (Aktas and Hasbenli 1994)	Long-winged bats <i>Miniopterus</i> spp.	Carrier of virus from family Rhabdoviridae (Aznar-Lopez et al., 2013)
<i>Phthiridium biarticulatum</i> (Hermann, 1804)	—	Algeria (Bendjedou et al., 2014)	Horseshoe microbats (Chiroptera: Rhinophidae), predominantly greater horseshoe bat <i>Rhinolophus ferrumequinum</i> (Schreber, 1774)	—
<i>Basilia daganaiae</i> Theodor and Moscona, 1954	—	Egypt (Theodor, 1956); Palestine (Theodor, 1967); Jordan (Benda et al., 2010) Cyprus (Bequaert, 1953; Theodor, 1967; Kock, 1974)	<i>P. kuhlii</i>	—
<i>Insecta: Siphonaptera: Ischnopsyllidae</i>				
<i>Ischnopsyllus octactenus</i> (Kolenati, 1856)	Rostov region, Astrakhan region (Zabashta et al., 2019; present article); Dagestan (present article)	Algeria (Bendjedou et al., 2013, 2017); Tunisia (Beaucouru and Kock, 1996); Morocco (Quetglas et al., 2014); Iran (Maleki-Ravasan et al., 2017) Turkey (Aktas 1987, 1990); Azerbaijan (Dubovchenko, 1968, 1969); Spain (Balearic Islands) (Quetglas et al., 2014)	Pipistrelles <i>Pipistrellus</i> spp.	Carry bacteria <i>Bartonella</i> sp. (Hornok et al., 2012)
<i>I. variabilis</i> (Wagner, 1898)*	Ulyanovsk region (present article)	—	Pipistrelles <i>Pipistrellus</i> spp.	—
<i>I. intermedius</i> (Rothschild, 1898)	—	Azerbaijan (Dubovchenko, 1965, 1969)	Different species of simple nosed bats family: pond bat <i>M. dasycneme</i> (Boie, 1825), common pipistrelle <i>P. pipistrellus</i> sensu stricto, serotine bat <i>Eptesicus serotinus</i> Schreber, 1774	—

Table 2. (Contd.)

Ectoparasite	Finds in Russia	Finds outside of Russia	Principal hosts	Medical importance
<i>I. dolosus</i> Dampf, 1912	—	Azerbaijan (Dubovchenko, 1969)	Some mouse-eared bat species	—
<i>I. consimilis</i> (Wahlgren, 1904)	—	Turkey (Smit, 1954), Palestine (Theodor and Moscona, 1954; Hopkins and Rothschild, 1956); Egypt (Lewis 1962; Hoogstraal and Traub, 1963; Haas and Tomich, 1973) Israel (Hopkins and Rothschild, 1956) Lebanon (Lewis, 1962) Libya (Hürka, 1982)	Pipistrelles <i>Pipistrellus</i> spp.	—
<i>Nycteridopsylla eusarca</i> Dampf, 1908	—	Czechoslovakia (Rosický, 1957)	Common noctule <i>N. noctula</i> (Schreber, 1774)	Carrier of bacteria <i>Rickettsia helvetica</i> Beati et al., 1993 (Hornok et al., 2012)
<i>N. pentactena</i> (Kolenati, 1856)	—	Azerbaijan (Ioff et al., 1965; Dubovchenko, 1968, 1969) Algeria (Bendjedou et al., 2017)	Long-eared bat <i>Plecotus</i> spp. and <i>Barbastella</i> spp.	—
<i>N. levantina</i> Jordan, 1942	—	Cyprus (Jordan, 1942)	Species is known on the basis of a single find	—
Insecta: Siphonaptera: Pulicidae				
<i>Xenopsylla cheopis</i> (Rothschild, 1903)	—	Palestine (Theodor and Moscona, 1954)	Rodents, predominantly rats, ( <i>Rattus</i> spp., <i>Nesokcia</i> spp.) and gerbils (Gerbillinae)	Carry bacteria <i>Yersinia pestis</i> , causative agent of endemic typhus <i>Rickettsia typhi</i> (Wolbach and Todd 1920) Philip 1943 (Shrewsbury, 2005; Christou et al., 2010)
<i>Leptopsylla sengis</i> (Schoenherr, 1911)	—	Palestine (Theodor and Moscona, 1954)	Rats <i>Rattus</i> spp.	Carry bacteria <i>Rickettsia typhi</i> (Christou et al., 2010)

Table 2. (Contd.)

Ectoparasite	Finds in Russia	Finds outside of Russia	Principal hosts	Medical importance
Insecta: Hemiptera: Cimicidae				
<i>Cimex pipistrelli</i> (Jenyns, 1839)/ <i>Cimex ex gr. pipistrelli</i>	Rostov region (Zabashta et al., 2019)	Algeria (Bendjedou et al., 2013, 2017)	Pipistrelles <i>Pipistrellus</i> spp.	Carrier of causative agents of tularemia <i>Francisella tularensis</i> (McCoy and Chapin 1912) Dorofe'ev 1947 (Zabashta et al., 2019)
<i>Cimex lectularius</i> Linnaeus, 1758	—	Lybia (Hufnagl, 1972); Iraq (Abul-Hab and Shihab, 1990)	Homeothermic animals, including humans	Carry bacteria <i>Francisella tularensis</i> ; causative agents of brucellosis <i>Bruccella</i> sp., hepatitis B virus, tuberculosis <i>Mycobacterium tuberculosis</i> (Zopf 1883) Lehmann and Neumann 1896, salmonellosis <i>Salmonella typhi</i> , and anthrax <i>Bacillus anthracis</i> Cohn 1872 (Shestopalov et al., 2012)
<i>Cacodmus vicinus</i> Horváth, 1934	—	Turkey (Aktas and Kiyak 1990); Cyprus (Quetglas et al., 2012); Syria (Quetglas et al., 2012); Lebanon (Usinger, 1966); Israel (Quetglas et al., 2012); Jordan (Usinger, 1966); Benda et al., 2010; Quetglas et al., 2012); Algeria (Bendjedou et al., 2017)	<i>P. kuhlii</i>	—
		Egypt (Usinger, 1966; Quetglas et al., 2012); Lybia (Hürka, 1982); Tunisia (Usinger, 1966); Chad (Péricart, 1996)		

\* Host-parasite association recorded for the first time. The table did not include finds of *Seatonysus musculi* (Schrank, 1803) and *Seatonysus murinus* Lucas, 1840 in Azerbaijan (Dubovchenko, 1968), inasmuch as currently these taxa are not considered valid.

**Table 3.** Pathogens isolated from the Kuhl's pipistrelle

Pathogen	Territory	Source	Diseases caused
Apicomplexa: Eimeriidae			
<i>Eimeria pipistrellus</i>	Saudi Arabia	Alyousif et al., 1999	Coccidiosis (eimeriosis)
<i>Eimeria chiropteri</i>		Alyousif, 1999a	
<i>Eimeria kuhliensis</i>		Alyousif, 1999b	
Bacteria: Spirochaetaceae			
<i>Borrelia</i> sp.	Rostov region, Russia	Zabashta et al., 2019	Lyme disease
<i>Borellia afzelii</i>			
<i>Borrelia burgdorferi</i> s. l. genospecies <i>Borrelia afzelii</i>			
Anaplasmataceae			
<i>Ehrlichia</i> sp.	Rostov region, Russia	Zabashta et al., 2019	Ehrlichiosis
Francisellaceae			
<i>Francisella tularensis</i>	Rostov region, Russia	Zabashta et al., 2019	Tularemia
Virus			
Vapiro virus (VAPV) (Rhabdoviridae)	Italy	Lelli et al., 2018	—
Toscana virus (TOSV) (Bunyaviridae)		Verani et al., 1988	—
Alphacoronavirus (Coronaviridae)	Italy, Spain	Lelli et al., 2013 Falcón et al., 2011	—
Betacoronavirus (Coronaviridae)	Italy	Lelli et al., 2013	—
Orthoreovirus (Reoviridae)	Germany, Italy	Kohl et al., 2012 Lelli et al., 2012	—

individuals ranges from 56 to 100%. The next abundant species in our material flea *I. octactenus* is characterized by smaller infestation parameters (MI from 1 to 2, P from 18 to 100%). The remaining two species (tick *C. vespertilionis* and flea *I. variabilis*) are represented by isolated individuals.

Thus, according to our own and published data, 37 arthropods identified to species were recorded from the Kuhl's pipistrelle (mites, ticks, and insects), among which 36 are parasites (Table 2). In addition, 13 pathogenic microorganisms were isolated from this species over the past 20 years: three protozoan species (coccidia) of genus *Eimeria*, three groups of *Borrelia* (*Borrelia burgdorferi* s. l., *Borrelia afzelii*, and *Borrelia* sp.), *Ehrlichia* sp., *Francisella tularensis*, and five viruses belonging to four families (Rhabdoviridae, Bunyaviridae, Coronaviridae, and Reoviridae) (Table 3).

In faunistic terms, arthropods are represented by Acari, including 16 species from eight families of three orders, and insects, including 18 species from four families of three orders.

Apparently, only some of the arthropod finds are due to parasitism. Thus, nonparasitic *Kleemannia* sp. appears to be an accidental find, because these mites are associated with bees. A number of finds are artifacts (mites *Laelaps algericus*, *Dermanyssus gallinae*,

*Spinturnix psi*, *Meristaspis lateralis*, *Parasteatonyssus ca. nyctinomi*, and *Hyalomma dromedarii*; bat flies *Nycteribia schmidlii*, *Phthiridium biarticulatum*, *Penicillidia conspicua*; fleas *Xenopsylla cheopis* and *Leptopsyllea sengis*). At the same time, these finds, in particular, characterize active contacts of the Kuhl's pipistrelle with other bat species, as well as vertebrates from other classes (birds) and orders (primarily, domestic animals, such as rats and mice). In this case, the Kuhl's pipistrelle acts as a primary host (or one of these hosts) only for 11 species of ectoparasites (five Acari and six insect species), the most abundant and common among which (Acari *St. periblepharus* and *C. vespertilionis*, fleas *I. octactenus* and *I. consimilis*, bat flies *B. daganaiae*, and bugs *Cacodmus vicinus*) appear to form a core of ectoparasitic *P. kuhlii* fauna.

Data on parasites of this species in Russia is on a smaller scale. As few as six arthropod species (three Acari and three insect species) and three taxa of pathogens were similarly recorded by us earlier, which can be attributed to insufficient knowledge about this host. Interestingly, all ectoparasitic species recorded in Russia are common to *Pipistrellus* spp. without being exclusively specific to *P. kuhlii*. As range of the bats expands northward, the core of their ectoparasitic fauna can be assumed to undergo a series of changes in

that, in particular, highly host-specific species are lost. The latter include ectoparasitic insects typical to the Kuhl's pipistrelle, such as bat fly *Basilia daganaiae*, flea *Ischnopsyllus consimilis*, and bug *C. vicinus*, the finds of which outside of the Mediterranean are unknown. All of the three listed species are transient ectoparasites sensitive, among other factors, to environmental parameters. Therefore, lack of the ecologically adequate conditions in their shelters and, primarily, not sufficiently high temperature can explain allocation (confinedness) of these insects to the ancestral range of *P. kuhlii*. In actual fact, outside of the Mediterranean, the core of ectoparasitic fauna of this host is represented only by pipistrelle genus-specific parasites (Acari *St. periblepharus* and *C. vespertilionis* and flea *I. octactenus*). Our finding from the Kuhl's pipistrelle of flea *I. variabilis* common to temperate belt of Europe, stretching from the Atlantic region to Urals, and not seen within the ancestral range of this host suggests that *P. kuhlii* can enter a spectrum of hosts of extrinsic parasites from allied bat species.

Of particular concern are the multiple finds (including accidental encounters) from the Kuhl's pipistrelle of bat flies, which are vectors of various diseases, such as *C. vespertilionis*, *L. algericus*, *D. gallinae*, *X. cheopis*, and *Lept. sengis*, as well as data with respect to the its isolated pathogens. Presumably, *P. kuhlii* is involved in transmission of a number of bacteria harmful to humans, e.g., *Borrelia*, *Ehrlichia*, and tularemia agent, and coronaviruses, for which a possibility of transmission to humans was confirmed by the 2020 pandemic (Table 3). Along with this, it has acquired a status of abundant species in some areas, after having become established in, e.g., many cities of European Russia. Additionally, as a bat, the Kuhl's pipistrelle is characterized by all traits common to this group, such as a capacity to cover long distances (can potentially spread pathogens over considerable area) and active use of anthropogenic structures for day roosts and hibernation roosts; in other words, it has contacts with humans and synanthropic species, including the ones involved in circulation of the natural focus infections, such as rats and house mice, and has the possibility to exchange pathogens. Altogether this determines urgency for the further in-depth inquiry into the ecology and medical importance of the Kuhl's pipistrelle and associated parasites in Russia.

#### ACKNOWLEDGMENTS

We thank the administration of the Astrakhan Nature Reserve and individually I.V. Sokolova, as well as the directorship and inspectors of the Dagestan State Nature Reserve and individually Deputy Director of Research G.S. Dzhamirzoev for their assistance in carrying out the work.

#### FUNDING

The study was performed within the framework of a state assignment of the Ministry of Science and Higher Education of the Russian Federation (project no. 0721-2020-0019).

#### COMPLIANCE WITH ETHICAL STANDARDS

*Conflict of interest.* The authors declare that they have no conflict of interest.

*Statement on the welfare of animals.* All experimental protocols were implemented in compliance with the guiding principles of the European Union for the care and use of animals (86/609/CEE) and in compliance with rules approved by the instruction of the Presidium of the USSR Academy of Sciences dated April 2, 1980, no. 12000-496, and the order of the Ministry of Higher Educational Institutions of the Soviet Union dated September 13, 1984, no. 22. Every effort was made to use the minimum number of animals necessary to obtain reliable scientific data.

#### REFERENCES

- Abd El-Baky, S.M.M., Prevalence of external parasites in the south eastern desert of Egypt, *J. Egypt. Soc. Parasitol.*, 2001, vol. 31, pp. 223–232.
- Abul-Hab, J. and Shihab, B.A., Ectoparasites of some bats from Iraq, *Bull. Iraq Nat. Hist. Mus.*, 1990, vol. 8, no. 2, pp. 59–64.
- Aktaş, M. and Hasbenli, A., Bat flies of eastern Turkey (the east of Samsun-İskenderun line) (Diptera: Nycteribidae), *Gazi Univ. Fen Bilimleri Enstitüsü Derg.*, 1994, no. 7, pp. 48–51.
- Aktaş, M. and Kiyak, S., A new record of *Cacodromus vicinus* Horváth to the fauna of Turkey (Hemiptera: Cimicidae), *Cent. Entomol. Stud. Ankara, Misc. Pap.*, 1990, no. 7, pp. 1–2.
- Aktaş, M., Bat fleas of Eastern Turkey (the east of Samsun-İskenderun line), *Doga – Turk. J. Zool.*, 1987, vol. 11, pp. 111–118.
- Aktaş, M., Bat fleas of western Turkey (west of the line between Samsun and İskenderun), *Doga – Turk. J. Zool.*, 1990, vol. 14, pp. 107–112.
- Al'khovskii, S.V., L'vov, D.K., Shchelkanov, M.Yu., Shchetinin, A.M., and Deryabin, P.G., Taxonomy of the Issyk-Kul virus (Issyk-Kul, ISKV; Bunyaviridae, Nairovirus), the causative agent of Issyk-Kul fever, isolated from bats (Vespertilionidae) and ticks Argas (Carrios) vespertilionis Latreille, 1796, *Vopr. Virusol.*, 2013, vol. 58, no. 5, pp. 11–15.
- Alyousif, M.S., Al-Dakhil, M., and Al-Shawa, Y.R., *Eimeria pipistrellus* n. sp. from *Pipistrellus kuhlii* (Chiroptera: Vespertilionidae) in Saudi Arabia, *Korean J. Parasitol.*, 1999, vol. 37, no. 1, pp. 1–4.
- Alyousif, M.S., *Eimeria chiropteri* n. sp. (Apicomplexa: Eimeriidae) from Saudi Arabian bat *Pipistrellus kuhlii* (Chiroptera: Vespertilionidae), *J. Egypt. Soc. Parasitol.*, 1999a, vol. 29, no. 1, pp. 275–280.
- Alyousif, M.S., *Eimeria kuhliensis* sp. n. (Apicomplexa: Eimeriidae) from pipistrelle bat, *Pipistrellus kuhlii*, *Acta Protozool.*, 1999b, no. 38, pp. 313–316.

- Anciaux de Faveaux, M., Distribution des chiroptères en Algérie, avec notes écologiques et parasitologiques, *Bull. Soc. Hist. Nat. Afr. Nord*, 1976, vol. 67, pp. 69–80.
- Ancillotto, L., Santini, L., Ranc, N., Maiorano, L., and Russo, D., Extraordinary range expansion in a common bat: The potential roles of climate change and urbanization, *Sci. Nat.*, 2016, no. 103, p. 15.
- Artaev, O.N and Smirnov, D.G., Bats (Chiroptera; Mammalia) of Mordovia: species composition and distribution features, *Nat. Conserv. Res., Zapov. Nauka*, 2016, vol. 1, no. 1, pp. 38–51.
- Arutyunyan, E.S. and Ogadzhanyan, A.M., Parasitic mites of family Spinturnicidae Oudemans, 1901 (Parasitiformes, Gamasoidea) of Armenian bats, *Biol. Zh. Arm.*, 1974, vol. 27, no. 4, pp. 72–81.
- Aznar-Lopez, C., Vazquez-Moron, S., Marston, D.A., Juste, J., Ibáñez, C., Berciano, J.M., Salsamendi, E., Aihartza, J., Banyard, A.C., McElhinney, L., Fooks, A.R., and Echevarria, J., Detection of rhabdovirus viral RNA in oropharyngeal swabs and ectoparasites of Spanish bats, *J. Gen. Virol.*, 2013, vol. 94, pp. 69–75.
- Balashov, Yu.S., *Parazitizm kleshchei i nasekomykh na nazemnykh pozvonochnykh* (Parasitism of Mites and Insects on Terrestrial Vertebrates), St. Petersburg: Nauka, 2009.
- Bauer, K., Ausbreitung der Weißrandfledermaus *Pipistrellus kuhlii* (Kuhl, 1819) in Österreich, *Mitt. Landesmus. Joan. Zool.*, 1996, no. 50, pp. 17–24.
- Beaucournu, J.-C. and Kock, D., Notes on the Ischnopsyllinae of the African continent, III. Additions to the distribution of species (Insecta: Siphonaptera: Ischnopsyllidae), *Senckenbergiana Biol.*, 1996, no. 75, pp. 163–169.
- Benda, P., Faizoláhi, K., Andreas, M., Obuch, J., Reiter, A., Ševčík, M., Uhrin, M., Vallo, P., and Ashrafi, S., Bats (Mammalia: Chiroptera) of the Eastern Mediterranean and Middle East. Part 10. Bat fauna of Iran, *Acta Soc. Zool. Bohemicae*, 2012, vol. 76, nos. 3–4, pp. 163–582.
- Benda, P., Lučan, R.K., Obuch, J., Reiter, A., Andreas, M., Backor, P., Bohnenstengel, T., Eid, E.K., Ševčík, M., Vallo, P., and Amr, Z.S., Bats (Mammalia: Chiroptera) of the Eastern Mediterranean and Middle East. Part 8. Bats of Jordan: fauna, ecology, echolocation, ectoparasites, *Acta Soc. Zool. Bohemicae*, 2010, vol. 74, pp. 185–353.
- Benda, P., Spitzerberger, F., Hanák, V., Andreas, M., Reiter, A., Ševčík, M., Šmid, J., and Uhrin, M., Bats (Mammalia: Chiroptera) of the Eastern Mediterranean and Middle East. Part 11. On the bat fauna of Libya II, *Acta Soc. Zool. Bohemicae*, 2014, vol. 78, pp. 1–162.
- Bendjedou, M.L., Bitam, I., Abiad, A., Bouslama, Z., and Amr, Z., New records of arthropod ectoparasites of bats from North–Eastern Algeria, *Jordan J. Biol. Sci.*, 2013, vol. 6, no. 4, pp. 324–327.
- Bendjedou, M.L., Loumassine, H.A., Scheffler, I., Bouslama, Z., and Amr, Z., Bat ectoparasites (Nycteribiidae, Streblidae, Siphonaptera, Heteroptera, Mesostigmata, Argasidae, and Ixodidae) from Algeria, *J. Vector Ecol.*, 2017, vol. 42, no. 1, pp. 13–23.
- Bequaert, J.G., Exploration du Parc National Albert. Mission G.F. de Witte (1933–1935). Hippoboscidae and Nycteribiidae, *Fasc. Inst. Parcs Nat.*, 1953, vol. 79, no. 7, pp. 89–94.
- Bogdanowicz, W., *Pipistrellus kuhlii* (Kuhl, 1817) – Weissrandfledermaus, in *Handbuch der Saugetiere Europas*, vol. 4: *Fledertiere*, part II: *Chiroptera II*, Krapp, F., Ed., Wiebelsheim: Aula-Verlag, 2004, pp. 875–908.
- Ceeľuch, M. and Ševčík, M., First record of *Pipistrellus kuhlii* (Chiroptera) from Slovakia, *Biologia* (Bratislava), 2006, vol. 61, pp. 637–638.
- Christou, C., Psaroulaki, A., Antoniou, M., Toumazos, P., Ioannou, I., Mazeris, A., Chochlakis, D., and Tselenitis, Y., *Rickettsia typhi* and *Rickettsia felis* in *Xenopsylla cheopis* and *Leptopsylla segnis* parasitizing rats in Cyprus, *Am. J. Trop. Med. Hyg.*, 2010, no. 83, pp. 1301–1304.
- Danko, Š., Reprodukcia *Hypsugo savii* a *Pipistrellus kuhlii* na východnom Slovensku: Daňšedokazy o ichšírenina sever, *Vespertilio*, 2007, no. 11, pp. 13–24.
- Davygora, A.V., Il'in, V.Yu., Smirnov, D.G., and Shepelev, A.A., Modern eastern limits of distribution of *Pipistrellus kuhlii* in Russia, in *Sovremennye problemy zoologii filogeografii mlekopitayushchikh. Materialy konferentsii* (Current Problems of Zoo- and Phylogeoigraphy of Mammals. Conference Proceedings), Pensa: Tov. Nauchn. Izd. KMK, 2009.
- Demyanchik, V.V., Faunistic complexes of vertebrates in the settlements of Pripyat Polesie, in *Ratsional'noe ispol'zovanie poimennyykh zemel'* (Rational Use of Floodplain Lands), Minsk, 2013, pp. 98–102.
- Dubovchenko, T.A., Ectoparasites of Azerbaijan bats, *Extended Abstract of Cand. Sci. (Biol.) Dissertation*, Baku, 1968.
- Dubovchenko, T.A., Materials for the study of fleas in bats in Azerbaijan, in *Voprosy parazitologii AN Azerbaidzhanskoi SSR* (Problems of Parasitology, Academy of Sciences of the Azerbaijan SSR), Baku: Elm, 1969, pp. 236–240.
- Dubovchenko, T.A., New species of fleas for the fauna of Azerbaijan, *Materialy nauchnoi sessii entomologov Azerbaidzhana* (Proc. Sci. Session of Entomologists of Azerbaijan), Baku: Akad. Nauk Azerb. SSR, 1965, pp. 83–84.
- Dusbabek, F. and Arutunian, E.S., *Acanthophthirius kolentati*, a new species, and some new records of fur mites (Acarina: Myobiidae and Labidocarpidae) from Armenia, *Folia Parasitol.* (Prague), 1976, vol. 23, pp. 57–60.
- Falcón, A., Vázquez-Morón, S., Casas, I., Aznar, C., Ruiz, G., Pozo, F., Pérez-Breña, P., Juste, J., Ibáñez, C., Garín, I., et al., Detection of alpha and betacoronaviruses in multiple Iberian bat species, *Arch. Virol.*, 2011, no. 156, pp. 1883–1890.
- Feher, C.E., A feherszelű denever (*Pipistrellus kuhlii*) erső magyarországi adatai, *Deneverkutatas – Hung. Bat Res. News*, 1995, no. 1, pp. 16–17.
- Fiedler, W., Alder, H.U., and Wohland, P., Zwei neue nachweise der Weißrandfledermaus (*Pipistrellus kuhlii*) für Deutschland, *Z. Saugetierk.*, 1999, vol. 64, pp. 107–109.
- Filippova, N.A., Argasidae, in *Fauna SSSR: Paukoobraznye* (Fauna of the USSR: Arachnids), Moscow: Nauka, 1966, vol. 4, no. 4.
- Flochlay, A.S., Thomas, E., and Sparagano, O., Poultry red mite (*Dermanyssus gallinae*) infestation: a broad impact parasitological disease that still remains a significant

- cant challenge for the egg-laying industry in Europe, *Parasites Vectors*, 2017, vol. 10, p. 357.
- Gadzhiev, A.T. and Dubovchenko, T.A., Gamasid mites (Parasitiformes, Gamasoidea) of bats of Azerbaijan, *Zool. Zh.*, 1967, vol. 46, no. 11, pp. 1716–1719.
- Gardner, R.A. and Molyneux, D.H., *Polychromophilus murinus*: a malarial parasite of bats: life-history and ultrastructural studies, *Parasitology*, 1988, vol. 96, pp. 591–605.
- Gashchak, S.P., Vlashchenko, A.S., and Naglov, A.V., Results of studying the fauna and radioactive contamination of bats in the Chernobyl exclusion zone in 2007–2009, *Probl. Chornobil's'koj Zoni Vidchuzhennya*, 2009, no. 9, pp. 102–124.
- Gavris', G.G. and Kotserzhinskaya, I.M., The record of *Pipistrellus kuhlii* (Chiroptera, Vespertilionidae) on the northern border of the range in Ukraine (Shchorsky rayon, Chernihiv region), *Vestn. Zool.*, 2002, vol. 36, no. 6, p. 50.
- Godlevs'ka, L., Results of the work of the Kyiv Contact Center for Bats in 2012–2015, *Pratsi Teriol. Shk.*, 2015, vol. 13, pp. 11–19.
- Godlevsky, L., Tyshchenko, V., and Negoda, V., First records of *Pipistrellus kuhlii* from Kyiv, *Vestn. Zool.*, 2000, vol. 34, no. 3, p. 78.
- Haas, G.E. and Tomich, P.Q., The fleas (Siphonaptera) of Egypt. Early records from mammals, *J. Egypt. Publ. Health Assoc.*, 1973, no. 48, pp. 1–17.
- Haffner, M., Stutz, H.-P.B., and Zumsteg, M., First record of Swiss nursery colonies of *Pipistrellus kuhlii* (Natterer in Kuhl, 1819) (Mammalia: Chiroptera) north of the Alps, *Rev. Suisse Zool.*, 1991, vol. 98, pp. 702–703.
- Hamed, M.I., Zaitoun, A.M.A., El-Allawy, T.A.A., and Mourad, M.I., Investigation of *Theileria camelensis* in camels infested by *Hyalomma dromedarii* ticks in Upper Egypt, *J. Adv. Vet. Res.*, 2011, vol. 1, pp. 4–7.
- Hoogstraal, H. and Traub, R., The fleas (Siphonaptera) of Egypt. Host-parasite relationships of Insectivora and Chiroptera, *J. Egypt. Publ. Health Assoc.*, 1963, vol. 38, pp. 111–130.
- Hoogstraal, H., A preliminary, annotated list of ticks (Ixodoidea) of the Anglo-Egyptian Sudan, *J. Parasitol.*, 1954, vol. 40, pp. 304–310.
- Hopkins, G.H.E. and Rothschild, M., *An Illustrated Catalogue of the Rothschild Collection of Fleas (Siphonaptera) in the British Museum (Natural History)*, vol. II: *Coptopsyllidae, Vermipsyllidae, Stephanocircidae, Ischnopsyllidae, Hypsophthalmidae and Xiphopsyllidae*, London, 1956.
- Hornok, S., Kovács, R., Meli, M.L., Gönczi, E., Hofmann-Lehmann, R., Kontschán, J., Gyuranecz, M., Dán, A., and Molnár, V., First detection of Bartonellae in a broad range of bat ectoparasites, *Vet. Microbiol.*, 2012, vol. 159, nos. 3–4, pp. 541–543.
- Hornok, S., Szőke, K., Göröföl, T., Földvári, G., Tu, V.T., Takács, N., Jenő, K., Sándor, A.D., Estók, P., Epis, S., Boldogh, S.A., Kováts, D., and Wang, Y., Molecular investigations of the bat tick *Argas vespertilionis* (Ixodida: Argasidae) and *Babesia vesperuginis* (Apicomplexa: Piroplasmida) reflect “bat connection” between Central Europe and Central Asia, *Exp. Appl. Acarol.*, 2017, vol. 72, pp. 69–77.
- Hornok, S., Szőke, K., Kováts, D., Estók, P., Göröföl, T., Boldogh, S.A., Takács, N., Jenő, K., Földvári, G., Barti, L., Corduneanu, A., and Sándor, A.D., DNA of piroplasms of ruminants and dogs in Ixodid bat ticks, *PLoS One*, 2016, vol. 11, no. 12.
- Hufnagl, E., *Libyan Mammals*, Stoughton & Harrow: The Oleander Press, 1972.
- Hürka, K., On the insect bat ectoparasites of coastal Libya (Cimicidae, Nycteribiidae, Streblidae, Ischnopsyllidae), *Vestn. Cesk. Spol. Zool.*, 1982, vol. 46, pp. 85–91.
- Il'in, V.Yu., Bystrakova, N.V., Dobrolyubov, A.N., Er-makov, O.A., Zolina, N.F., Kurmaeva, N.M., Luk'yanov, S.B., Pavlova, S.V., Smirnov, D.G., and Titov, S.V., Abstract of the mammalian fauna of Penza region, *Izv. Penz. Gos. Pedagog. Univ. im. V. G. Belinskogo*, 2006, vol. 1, no. 5, pp. 73–89.
- Il'in, V.Yu., Dynamics of the ranges of three species of bats in the extreme southeast of Europe, *Plecotus et al.*, 2000, no. 3, pp. 43–49.
- Imaz, E., Aihartza, J.R., and Totorika, M.J., Ectoparasites on bats (Gamasida, Ixodida, Diptera) in Biscay (N Iberian peninsula), *Misc. Zool.*, 1999, vol. 22, pp. 21–30.
- International Catalogue of Arboviruses Including Certain Other Viruses of Vertebrates*, Karabatos, N., Ed., San Antonio, TX, Am. Soc. Trop. Med. Hyg., 1985.
- Ioff, I.G., Mikulin, M.A., and Skalon, O.I., *Opredelitel' blokh Srednei Azii i Kazakhstana* (Keys to the Fleas of Central Asia and Kazakhstan), Moscow: Meditsina, 1965.
- Jordan, K., On four new Palearctic bat-fleas in the British Museum collection, *Eos* (Madrid), 1942, vol. 18, pp. 243–250.
- Kapitonov, V.I. and Grigor'ev, A.K., On the record of Kuhl's pipistrelle (*Pipistrellus kuhlii*) in Udmurtia, *Tez. Dokl. 2-i Rossiiskoi univers. akademich. nauch. prakt. konf.* (Reports of the 2nd Russian University Academic Scientific Practical Conf.), Izhevsk, 1995, part 2, pp. 40–41.
- Kedrov, B.M. and Sheshurak, P.N., The first find of the Mediterranean bat *Pipistrellus kuhlii* (Chiroptera, Vespertilionidae) in Chernigov region (Ukraine), *Vestn. Zool.*, 1999, vol. 33, no. 3, p. 66.
- Kock, D., Pupipare Dipteren von Säugetieren des nordöstlichen Mittelmeerraumens (Ins.: Diptera), *Senckenberg. Biol.*, 1974, vol. 17, pp. 87–104.
- Kohl, C., Lesnik, R., Brinkmann, A., Ebinger, A., Radonić, A., Nitsche, A., Mühlendorfer, K., Wibbelt, G., and Kurth, A., Isolation and characterization of three mammalian orthoreoviruses from European bats, *PLoS One*, 2012, vol. 7, no. 8, pp. 106–149.
- Komarov, Yu.E. and Kuchiev, I.T., Distribution of summer colonies of some bats in Ossetia, in *Ekol. gorn. mlekopitayushchikh: Inf. materialy* (Ecology of Mountaneous Mammals: Information Materials), Sverdlovsk, 1982, pp. 56–57.
- Korine, C., Pilosof, S., Gross, A., Morales-Malacara, J.B., and Krasnov, B.R., The effect of water contamination and host-related factors on ectoparasite load in an insectivorous bat, *Parasitol. Res.*, 2017, vol. 116, no. 9, pp. 2517–2526.
- Kuzyakin, A.P., *Letuchie myshi* (Bats), Moscow: Sov. Nauka, 1950.
- Lada, G.A., The record of Kuhl's pipistrelle in Tambov region, *Zool. Zh.*, 2010, vol. 89, no. 7, pp. 888–890.
- Lelli, D., Moreno, A., Lavazza, A., Bresaola, M., Canelli, E., Boniotti, M.B., and Cordioli, P., Identification of mammalian orthoreovirus type 3 in Italian bats, *Zoonoses Publ. Health*, 2012, vol. 60, pp. 84–92.

- Lelli, D., Papetti, A., Sabelli, C., Rosti, E., Moreno, A., and Boniotti, M.B., Detection of coronaviruses in bats of various species in Italy, *Viruses*, 2013, vol. 5, pp. 2679–2689.
- Lelli, D., Prosperi, A., Moreno, A., Chiapponi, C., Gibellini, A.M., De Benedictis, P., Leopardi, S., Sozzi, E., and Lavazza, A., Isolation of a novel Rhabdovirus from an insectivorous bat (*Pipistrellus kuhlii*) in Italy, *Virol. J.*, 2018, vol. 15, no. 1, p. 37.
- Lewis, R.E., A preliminary list of the fleas of Lebanon, Proceedings of the Royal Entomological Society of London, *Gen. Entomol., Ser. A*, 1962, vol. 37, pp. 49–60.
- Lindeman, G.V. and Subbotin, A.E., On the passage of bats in clay semideserts of the Trans-Volga region, *Byull. MOIP, Otd. Biol.*, 1983, vol. 88, no. 4, pp. 88–90.
- Lvov, D.K., Karas, F.R., Timofeev, E.M., Tsyrkin, Y.M., Vargina, S.G., Veselovskaya, O.V., Osipova, N.Z., Grebenyuk, Y.I., Gromashevski, V.L., Steblyanko, S.N., and Fomina, K.B., "Issyk-Kul" virus, a new arbovirus isolated from bats and *Argas (Carios) vespertilionis* (Latr., 1802) in the Kirghiz S.S.R., *Brief Rep., Arch. Ges. Virusforsch.*, 1973, vol. 42, no. 2, pp. 207–209.
- Maleki-Ravasan, N., Soljhony-Fard, S., Beaucournu, J.-C., Laudisoit, A., and Mostafavi, E., The fleas (Siphonaptera) in Iran: Diversity, host range, and medical importance, *PLoS Neglected Trop. Dis.*, 2017, vol. 11, no. 1. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5249249>. Accessed August 3, 2020.
- Meschede, A., Schlapp, G., and Weid, R., Erstfund einer Weißbrandfledermaus (*Pipistrellus kuhlii* Kuhl, 1819) in Bayern, *Nyctalus (N. F.)*, 1998, vol. 6, pp. 547–550.
- Moratelli, R. and Calisher, C.H., Bats and zoonotic viruses: can we confidently link bats with emerging deadly viruses?, *Mem. Inst. Oswaldo Cruz*, 2015, vol. 110, no. 1, pp. 1–22.
- Oba, M., Omatsu, T., Takano, A., Fujita, H., Sato, K., Nakamoto, A., Takahashi, M., Takada, N., Kawabata, H., Ando, S., and Mizutani, T., A novel Bunyavirus from the soft tick, *Argas vespertilionis*, in Japan, *J. Vet. Med. Sci.*, 2016, vol. 78, no. 3, pp. 443–445.
- Ogadzhanyan, A.M. and Arutyunyan, E.S., Ticks of the family Macropyssidae Oudemans, 1936 (Parasitiformes, Gamasoidea) parasitizing on bats of Armenia, *Biol. Zh. Arm.*, 1974, vol. 27, no. 10, pp. 75–82.
- Orlova, M.V. and Kononova, Yu.V., The causative agents of infectious diseases associated with specific ectoparasites of bats (Chiroptera) (review of literature data), *Parazitologiya*, 2018, vol. 52, no. 2, pp. 137–153.
- Orlova, M.V., Stanyukovich, M.K., and Orlov, O.L., *Gamasid Mites (Mesostigmata: Gamasina) Parasitizing Bats (Chiroptera: Rhinolophidae, Vespertilionidae, Molossidae) of Palaearctic Boreal Zone (Russia and Adjacent Countries)*, Tomsk: TSU Publ. House, 2016.
- Péricart, J., Family Cimicidae Latreille, 1802—bed-bugs, in *Catalogue of the Heteroptera of the Palearctic Region*, Aukema, B. and Rieger, C., Eds., Amsterdam: Neth. Entomol. Soc., 1996, pp. 141–144.
- Poluda, A. and Zagorodnyuk, I., Migrations of bats at the ornithological station "Lebedivka," in *Migraciinii status kazhaniv v Ukraini* (Migration Status of Bats in Ukraine), Kyiv, 2001, pp. 98–101.
- Quetglas, J., Balvín, O., Lučan, R.K., and Benda, P., First records of the bat bug *Cacodmus vicinus* (Heteroptera: Cimidae) from Europe and further data on its distribution, *Vespertilio*, 2012, vol. 16, pp. 243–248.
- Quetglas, J., Nogueras, J., Ibáñez, C., and Beaucournu, J.-C., Presencia en la Península Ibérica de una pulga africana de murciélagos: *Rhinolophopsylla unipectinata arabs* (Siphonaptera: Ischnopsyllidae) y otras nuevas citas de pulgas de murciélagos para España y Marruecos, *Galemys*, 2014, vol. 26, pp. 41–47.
- Racey, P.A., Reproductive assessment of bats, in *Ecological and Behavioural Methods for the Study of Bats*, Kunz, T.H. and Parsons, S., Eds., Baltimore: Johns Hopkins Univ. Press, 2009, pp. 249–264.
- Radovsky, F., *The Macropyssidae and Laelapidae (Acarina: Mesostigmata) Parasitic on Bats*, Berkeley: Univ. Ca., 1967.
- Reeves, W.K., Dowling, A.P., and Dasch, G.A., Rickettsial agents from parasitic Dermanysoidea (Acari: Mesostigmata), *Exp. Appl. Acarol.*, 2006, vol. 38, nos. 2–3, pp. 181–188.
- Reiter, A., Benda, P., and Hotovy, J., First record of the Kuhl's pipistrelle, *Pipistrellus kuhlii* (Kuhl, 1817), in the Czech Republic, *Lynx*, 2007, vol. 38, pp. 47–54.
- Rosický, B., Blechy – Aphaniptera, in *Fauna ČSR*, Praha: Nakladatelství Česk. Akad. Věd, 1957, vol. 10.
- Sachnowicz, K., Piskorski, M., and Tereba, A., Systematics and taxonomy of *Pipistrellus kuhlii* (Kuhl, 1817) in central Europe and the Balkans, *Zootaxa*, 2017, vol. 4306, no. 1, pp. 53–66.
- Sachnowicz, K., Wower, F., and Bashta, A.-T., Further range extension of *Pipistrellus kuhlii* (Kuhl, 1817) in central and eastern Europe, *Acta Chiropterol.*, 2006, vol. 8, no. 2, pp. 543–548.
- Schountz, T., Virology and immunology of bats, in *Bat Evolution, Ecology, and Conservation*, Adams, R.A. and Pedersen, S.C., Eds., New York: Springer Sci., 2013, pp. 393–412.
- Shaimardanov, R.T., New records (1985–1993) of species of bats (Chiroptera) that are rare and little known for the fauna of Kazakhstan, *Plecotus et al.*, 2001, no. 4, pp. 82–83.
- Sharifi, M., Mozafari, F., Taghinezhad, N., and Javanbakht, H., Variation in ectoparasite load reflects life history traits in the lesser mouse-eared bat *Myotis blythii* (Chiroptera: Vespertilionidae) in western Iran, *J. Parasitol.*, 2008, vol. 94, no. 3, pp. 622–625.
- Shepelev, A.A., Il'in, V.Yu., and Smirnov, D.G., Rare species of bats in Penza and its environs, in *Bioraznoobrazie: problemy i perspektivy sokhraneniya. Materialy mezhdunarodnoi nauchnoi konferentsii, posvyashchennoi 135-letiyu so dnya rozhdeniya I.I. Sprygina* (Biodiversity: Problems and Prospects for Conservation. Proc. Int. Sci. Conf. Dedicated to the 135th Anniversary of the Birth of I.I. Sprygin), Pensa, 2008, part 2, pp. 304–305.
- Shestopalov, N.V., Roslavtseva, S.A., Alekseev, M.A., Eremina, O.Yu., Tsarenko, V.A., and Osipova, N.Z., Epidemiological significance of bed bugs and measures to combat them. Information letter, *Dezinfektsionnoe Delo*, 2012, vol. 81, no. 3, pp. 42–46.
- Shpak, A. and Larchenko, A., Range expansion of Kuhl's pipistrelle (*Pipistrellus kuhlii*) into Belarus, *Proc. Theriol. Sch.*, 2016, vol. 14, pp. 99–102.
- Shrewsbury, J.F.D., *A History of Bubonic Plague in the British Isles*, Cambridge: Cambridge Univ. Press, 2005.

- Sitnikova, E.F., Kruskop, S.V., and Mishta, A.V., Materials on the bats fauna of Bryansk region, *Plecotus et al.*, 2009, nos. 11–12, pp. 32–49.
- Smirnov, D.G. and Vekhnik, V.P., On the current distribution of *Pipistrellus kuhlii* (Chiroptera: Vespertilionidae) in the Volga region, *Povolzh. Ekol. Zh.*, 2011, no. 2, pp. 193–202.
- Smirnov, D.G., Vekhnik, V.P., Kurmaeva, N.M., and Shepelev, A.A., New information on the distribution of *Pipistrellus kuhlii* in the Volga region, *Teriofauna Rossii i sopredel'nykh territorii (VIII s"ezd VTO). Materialy mezhdunarodnogo soveshchaniya* (Theriofauna of Russia and Adjacent Territories (VIII Congress of the All-Russian Theriological Society). Materials of the International Meeting), Moscow, 2007, p. 463.
- Smit, F.G.A.M., New bat-fleas (Siphonaptera: Ischnopsyllidae), *Parasitology*, 1954, vol. 44, pp. 144–156.
- Socolovschi, C., et al., *Borrelia*, *Rickettsia*, and *Ehrlichia* species in bat ticks, France, 2010, *Emerging Infect. Dis.*, 2012, vol. 18, no. 12, pp. 1966–1975.
- Spitzenberger, F. and Bauer, K., Weißbrandfledermaus *Pipistrellus kuhlii* (Kuhl, 1817), in *Die Säugetierfauna Österreichs. Bundesministerium für Land- und Forstwirtschaft Umwelt und Wasserwirtschaft*, Spitzenberger, F., Ed., Graz, 2001, pp. 245–248.
- Stanyukovich, M.K., Keys to the gamasid mites (Acari: Parasitiformes, Mesostigmata, Macronyssoidae et Laelaptoidea) parasiting bats (Mammalia, Chiroptera) from Russia and adjacent countries, *Rudolstädter Naturhist. Schr.*, 1997, vol. 7, pp. 13–46.
- Stekolnikov, A.A. and Quetglas, J., Bat-infesting chiggers (Acariformes: Trombiculidae) of the Balearic Islands and new data on the genus *Trisetica* Traub et Evans, 1950, *Folia Parasitol.*, 2019, vol. 66.
- Strelkov, P.P. and Il'in, V.Yu., Bats (Chiroptera, Vespertilionidae) in the south of the Middle and Lower Volga regions, *Tr. Zool. Inst. Akad. Nauk SSSR*, 1990, vol. 225, pp. 42–167.
- Strelkov, P.P., Expansion of ranges by Palaearctic bats (Chiroptera, Mammalia) as an example of invasion in anthropogenic biotopes, in *Biologicheskie invazii v vodnykh i nazemnykh ekosistemakh* (Biological Invasions in Aquatic and Terrestrial Ecosystems), Moscow: Tov. Nauchn. Izd. KMK, 2004, pp. 202–207.
- Strelkov, P.P., Order Chiroptera, Bats, in *Mlekopitayushchie Fauny SSSR. Ch. 1* (Mammals of the Fauna of the USSR. Part 1), 1963, pp. 134–218.
- Strelkov, P.P., Unkurova, V.I., and Medvedeva, G.A., New data on Kuhl's pipistrelle (*Pipistrellus kuhlii*) and dynamics of its range in the USSR, *Zool. Zh.*, 1985, vol. 64, no. 1, pp. 87–97.
- Theodor, O. and Moscona, A., On bat parasites in Palestine. I. Nycteribiidae, Streblidae, Hemiptera, Siphonaptera, *Parasitology*, 1954, vol. 44, pp. 157–245.
- Theodor, O., *An Illustrated Catalogue of the Rothschild Collection of Nycteribiidae in the British Museum (Natural History), with Keys and Short Descriptions for the Identification of Subfamilies, Genera, Species and Subspecies*, London: Br. Mus. (Nat. Hist.) Publ., 1967.
- Theodor, O., On the genus *Tripselia* and the group of *Basilia bathybothyra* (Nycteribiidae, Diptera), *Parasitology*, 1956, vol. 46, nos 3–4, pp. 353–394.
- Usinger, R.L., *Monograph of Cicadidae*, Washington, DC: Entomol. Soc. Am., 1966.
- Verani, P., Ciufolini, M.G., Caciolli, S., Renzi, A., Nicoletti, L., Sabatinelli, G., Bartolozzi, D., Volpi, G., Amaducci, L., and Coluzzi, M., Ecology of viruses isolated from sand flies in Italy and characterized of a new Phlebovirus (Arabia virus), *Am. J. Trop. Med. Hyg.*, 1988, vol. 38, pp. 433–439.
- Vereshchagin, N.K., *Mlekopitayushchie Kavkaza* (Mammals of the Caucasus), Moscow: Akad. Nauk SSSR, 1959.
- Vernier, E. and Bogdanowicz, W., *Pipistrellus kuhlii* (Kuhl, 1817), in *The Atlas of European Mammals*, Mitchell-Jones, A.J., Amori, G., Bogdanowicz, W., Krystufek, B., Reinjders, P.J.H., Spitzenberger, F., Stubbe, M., Thisen, J.B.M., Vohralík, V., and Zima, J., Eds., London: Acad. Press, 1999, pp. 120–121.
- Whitaker, Jr., J.O., Collecting and preserving ectoparasites for ecological study, in *Ecological and Behavioral Methods for the Study of Bats*, Washington: Smithsonian Inst. Press, 1988, pp. 459–474.
- Yarmysh, N.N. and Kazakov, B.A., Records of rare species in the Ciscaucasia, in *Redkie vidy mlekopitayushchikh i ikh okhrana: Materialy II Vsesoyuz. soveshch.* (Rare Species of Mammals and Their Protection: Proc. II All-Union Meeting), Moscow: Nauka, 1977, pp. 65–66.
- Yarmysh, N.N., Kazakov, B.A., Sonina, I.Yu., and Usvaiskaya, A.Kh., New records of bats in the North Caucasus, in *Rukokrylye (Chiroptera)* (Bats. (Chiroptera)), Moscow: Nauka, 1980, pp. 72–77.
- Zabashta, M.V., Orlova, M.V., Pichurina, N.L., Khametova, A.P., Romanova, L.V., Borodina, T.N., and Zabashta, A.V., Participation of bats (Chiroptera, Mammalia) and their ectoparasites in the circulation of pathogens of natural focal infections in southern Russia, *Parazitologiya*, 2019, vol. 53, no. 1, pp. 3–13.
- Zagorodnyuk, I.V. and Reznik, O.S., Expansion of the dark-headed form into the range of *Garrulus glandarius* in Donbass, *Berkut*, 2007, vol. 16, no. 1, pp. 103–109.
- Zav'yalov, E.V. and Shlyakhtin, G.V., Dynamics of the northern border of the range of the Mediterranean bat in the European part of Russia, in *Tez. dokl. VI s"ezda Teriologicheskogo ob-va v In-te problem ekologii i evolyutsii im. A.N. Severtsova RAN* (Reports of the VI Congress of the Teriological Society in Severtsov Institute of Problems of Ecology and Evolution, Russian Academy of Sciences), Moscow, 1999, p. 92.
- Zemskaya, A.A., *Paraziticheskie gamazovye kleshchi i ikh meditsinskoe znachenie* (Parasitic Gamasid Mites and Their Medical Significance), Moscow: Meditsina, 1973.
- Zolina, N.F., Shepelev, A.A., Smirnov, D.G., and Shepelev, A.A., Preliminary information about bats in Penza and its environs, *Povolzh. Ekol. Zh.*, 2007, no. 2, pp. 116–127.

*Translated by E. Kuznetsova*